Contents of Vol. 25, 1988

Baltisberger, M.: Atlas Florae Europaeae notes. 8. New nomenclatural	
combinations within Ranunculus polyanthemos	293-294
Czeczuga, B. & Skult, H.: Carotenoids in lichens of Southern Fin-	
land	229-232
Flatberg, K.I.: Taxonomy of Sphagnum annulatum and related spe-	
cies	303-350
Frahm, JP., Norris, D.H. & Koponen, T.: Bryophyte flora of the	505 550
Huon Peninsula, Papua New Guinea. XXV. Campylopodiella	
(Dicranaceae, Musci)	259-260
Gartmann, F.: Habitat-related differences between the vicarious fern	239-200
species Gympogornium dryontoric and C. rehardings term	261 274
species Gymnocarpium dryopteris and G. robertianum	261–274
Hedenäs, L.: The status of Orthothecium lapponicum and O. complana-	
tum (Musci, Plagiotheciaceae)	153–157
Heikkilä, R. & Lindholm, T.: Distribution and ecology of Sphagnum	
molle in Finland	11-19
Huovinen, K. & Ahti, T.: The composition and contents of aromatic	
lichen substances in Cladonia section Perviae	371-383
Jalas, J.: Atlas Florae Europaeae notes. 9–11. Ranunculaceae	295–299
Kanerva, T., Sarin, O. & Nuorteva, P.: Aluminium, iron, zinc, cad-	275 277
mium and mercury in some indicator plants growing in south	
Finnish forest areas with different degrees of damage	275–279
Vontamen V & Diel- C. Terrange GH.	213-219
Karttunen, K. & Bäck, S.: Taxonomy of Hymenodon (Musci, Rhizo-	00.0
goniaceae)	89–95
Koski, E., Venäläinen, M. & Nuorteva, P.: The influence of forest	
type, topographic location and season on the levels of Al, Fe, Zn,	
Cd and Hg in some plants in Southern Finland	365-370
Kožuharov, S. & Petrova, A.: Atlas Florae Europaeae notes. 12. News	
and adjustments concerning Bulgarian Ranunculaceae	389-390
Kullman, L.: Short-term dynamic approach to tree-limit and thermal cli-	
mate: evidence from Pinus sylvestris in the Swedish Scandes	219-227
Kuosa, H.: Some planktonic rock-pool algae from the Tvärminne archi-	219-221
	111–116
pelago, SW coast of Finland	111-110
Kuosa, H.: Estimating the proportion of active blue-green algal hetero-	
cysts in Baltic Sea samples by the TTC method	233–236
Lahti, T., Kurtto, A. & Väisänen, R.A.: Floristic composition and re-	
gional species richness of vascular plants in Finland	281-291
Lempiäinen, T.: Pflanzliche Makroreste von dem Innenhof des Qwen-	
selschen Anwesens in Turku, SW Finnland, vom 17.–19. Jh.	
(Abstract: Macrofossils from the Qwensel house in Turku, SW	
Finland, dated to the 17th, 18th and 19 centuries)	47-54
Leskinen, E. & Sarvala, J.: Community analysis of diatom colonization	17 51
on artificial substrata in a northern Baltic Sea archipelago: A com-	
	21 22
parison of methods	21–32
Nielsen, R.: Small green algae from brackish water in the Tvärminne	
area, southern Finland	237–257
Norris, D.H. & Koponen, T.: Bryophyte flora of the Huon Peninsula,	
Papua New Guinea. XXIV. Andreaeaceae, Archidiaceae, Seli-	
geriaceae, and Ditrichaceae (Musci)	165-177
Norris, D.H., Reese, W.D. & Koponen, T.: Bryophytes from Frieda	
River, East and West Sepik Provinces, Papua New Guinea. III.	
Syrrhopodon curticancellinatus, species nova (Calymperaceae)	149-151
Oksanen, J.: Impact of habitat, substrate and microsite classes on the	
eninhute vegetation. Interpretation using evaluratory and cononi	
epiphyte vegetation: Interpretation using exploratory and canoni-	50 71
cal correspondence analysis	59–71
Pietiläinen, P. & Lähdesmäki, P.: Effect of various concentrations of	
potassium nitrate and ammonium sulphate on nitrate reductase ac-	

	e roots and needles of Scots pine seedlings in N Fin-	201–206	
land	bryophyte flora of the Huon Peninsula, Papua New	201–200	
Guinea. XX	II. Lepicolea norrisii (Lepicoleaceae, Hepaticae)	55-57	
Piippo, S.: The	bryophyte flora of the Huon Peninsula, Papua New		
	XII. Targioniaceae, Wiesnerellaceae, Aytoniaceae and	97–107	
Piinno S · The	(Marchantiales, Hepaticae)bryophyte flora of the Huon Peninsula, Papua New	97-107	
Guinea. X	XIII. Treubiaceae, Allisoniaceae and Makinoaceae		
(Metzgeri	ales, Hepaticae)	159-164	
	dén, CJ., Salemink, C.A., Lewing, E., Rousi, A. &		
	interspecific hybridization in Papaver. 2. Alkaloid con-		
their inte	somniferum and species of the section Oxytona and rspecific hybrids	1-10	
Ravanko, O.: The	e chromosomes in three Chara species (Charophyta) in	1 10	
the SW a	rchipelago of Finland	85–88	
Ruokolahti, C. &	& Rönnberg, O.: Seasonal variation in chlorophyll a	205 200	
	Fucus vesiculosus in a northern Baltic archipelago w Taraxacum species from Albania. II	385–388 73–83	
Sorsa, P.: Pollen	morphology of Potamogeton and Groenlandia (Pota-	75-05	
mogetonac	eae) and its taxonomic significance	179-199	
Stenroos, S.: The	e family Cladoniaceae in Melanesia. 3. Cladonia sec-		
tions Held	opodium, Perviae and Cladonia	117–148	
	family Cladoniaceae in Melanesia. 4. The genera Claa, Calathaspis and Thysanothecium	207-217	
Tuomisto, H.: Us	se of Picea abies needles as indicators of air pollution:	207-217	
epicuticula	r wax morphology	351-364	
Vuorela, I. & Lempiäinen, T.: Archaeobotany of the site of the oldest			
cereal gran	n find in Finland	33–45	
Book Reviews:	Ahti, T.	58	
	Eurola, S.	384	
	Hæggström, CA.	84	
	Stenroos, S.	301, 302	
	Suominen, J.	20	
Correction	***************************************	109	
Numbers distributed			
Vol. 24(4):311–404			
Vol. 25(1):1–109			
Vol. 25(2):111–206			
Vol. 25(3):207-	302 2	26.X.1988	